

# Herbicide Programs for non-GMO Soybeans

Mark Loux  
**Horticulture and Crop Science**  
**The Ohio State University**  
<http://agcrops.osu.edu/weeds>

Bill Johnson  
 Glenn Nice  
**Purdue Extension Weed Science**  
[www.btny.purdue.edu/weedscience](http://www.btny.purdue.edu/weedscience)

**For Free Herbicide Labels Go to:**  
[www.cdms.net](http://www.cdms.net)  
 or  
[www.greenbook.net](http://www.greenbook.net)



**Purdue Extension**  
**Knowledge to Go**  
 1-888-EXT-INFO



## Essential components of non-GMO weed management

- Weed-free start at planting through use of tillage or preplant burndown herbicides
- Apply residual (PRE) herbicides in spring before soybeans emerge, to reduce weed populations, slow weed growth, and create flexibility in the postemergence application window
- Apply the first postemergence (POST) treatment when weeds are small - less than 4 to 6 inches
- Where needed to control late-emerging weeds, a second POST treatment should be applied about 3 weeks after the first POST treatment

## Spring preplant burndown programs

- Burndown should include either:
  - a. glyphosate (0.75 lb or higher) + 2,4-D ester (0.5 lb or higher), or
  - b. Ignite (29 to 36 oz) + 2,4-D ester (0.5 lb or higher) + metribuzin (0.38 lb ai), or
  - c. paraquat + 2,4-D ester (0.5 lb or higher) + metribuzin 75DF (6 oz or higher).
- Use of 2,4-D ester and soybean planting
  - a. 0.5 lb – 7 days before planting (any ester product)
  - b. 1.0 lb – 15 days before planting for Salvo, E99, Weedone 650 (30 days for other ester products)
- Apply when marestail are less than 4 inches tall – the smaller the better
- For most effective dandelion control, apply a mixture of glyphosate + 2,4-D ester + a residual product that contains chlorimuron or cloransulam.

## Use a broad-spectrum residual (PRE) herbicide

- residual should accomplish the following: 1) control lambsquarters and marestail, including ALS-resistant populations; 2) have activity on giant and common ragweed, including ALS-resistant populations where possible.
- Herbicide effectiveness ratings of 8 to 9 indicate “control”, and ratings of 6 or 7 indicate “has activity on”. More extensive ratings can be found in the “Weed Control Guide for Ohio and Indiana”.

	AMG	BNS	CRW	CRW ALSR	GRW	GRW ALSR	LBQ	MTL	MTL ALSR	PIG	VEL	WAH
<i>Auth Assist</i>	8	9	6	-	-	-	9	8	8	9	8	8
<i>Auth First/Sonic</i>	8	8	9	-	7	-	9	9	8	9	8	8
<i>Canopy/Cloak DF</i>	7	-	9	-	7	-	9	9	6	9	8	6
<i>Envive/ValorXLT</i>	8	9	9	7	7	-	9	9	8	9	8	8
<i>Gangster</i>	8	9	9	7	7	-	9	9	8	9	8	8
<i>Metribuzin</i>	-	-	7	7	-	-	9	8	8	9	7	8
<i>Python</i>	-	8	7	-	-	-	9	8	-	8	8	-
<i>Scepter</i>	7	9	8	-	7	-	9	6	-	9	7	-
<i>Valor</i>	7	9	7	7	-	-	9	8	8	9	7	8

Weed abbreviations: AMG = annual morningglory; BNS = black nightshade, CRW = common ragweed; GRW = giant ragweed, LBQ = lambsquarters, MTL = marestail, PIG = pigweed, VEL = velvetleaf, WAH = waterhemp;

## Suggested POST programs

- Apply first POST when weeds are less than 4 to 6 inches tall. Use spray volume of at least 15 gpa and nozzles that produce medium-sized droplets.
- Flexstar/Rhythm (1.3 - 1.5 pts/A) + grass herbicide (Select, Fusion, etc) + COC or MSO + AMS
  - a. can replace AMS with 28%, which may improve weed control but also increases crop injury
  - b. can add FirstRate, Classic or Synchrony to improve control of ALS-sensitive weeds
- Apply Cobra or Phoenix (10 oz/A) 3 weeks later if ragweed not dead, or to control late-emergers
  - a. can add low rate of grass herbicide (Select, Fusion, etc) for small late-emerging grass
- In fields without ALS-resistant weeds (be very sure about this before you go this route), it is possible to take the following approach in the first POST application: FirstRate (0.3 oz/A) or Classic (2/3 to 3/4 oz/A) + grass herbicide + COC or MSO + AMS. Flexstar/Rhythm (3/4 to 1 pt/A) can be added to improve control of ragweeds. In STS soybeans, a full rate of Synchrony XP can be used instead of FirstRate or Classic.

### Note on fall herbicide treatments

Residual herbicides are most effective and long-lasting when applied in the spring, not in the fall. Fall herbicide treatments should be used to manage winter annuals and dandelions, and should generally be followed by a spring preplant treatment that includes residual herbicides (in other words, don't substitute the fall treatment for a spring preplant treatment). Where a fall application is necessary, we suggest applying either: glyphosate + 2,4-D; or 2,4-D + a low rate of Canopy/Cloak EX/DF or other chlorimuron-containing product. This should be followed with a spring preplant application of a low rate of 2,4-D or glyphosate plus a residual herbicide. It is possible to follow fall application of a low rate of Canopy/Cloak EX/DF with another chlorimuron-containing herbicide in the spring, as long as the total amount of chlorimuron between fall and spring applications does not exceed maximum labeled rate for soil type.

## Approximate cost of herbicide programs for nonGMO soybeans

	ALS-resistant weeds	No ALS-resistant weeds
Preplant burndown	\$8 to \$12	\$8 to \$12
Residual (preemergence)	\$12 to \$16	\$12 to \$16
Postemergence (one app)	\$28 to \$44	\$17 to \$41
Total cost	\$48 to \$72	\$37 to \$69

In fields where a second POST treatment is needed, the total cost will increase by approximately \$15. Costs shown here do not include the cost of application, which typically ranges from \$4 to \$8 per acre.

Information listed here is based on research and outreach Extension programming at Purdue University, Ohio State University, and elsewhere. The use of trade names is for clarity to readers of this publication and does not imply endorsement of a particular brand nor does exclusion imply non-approval. Always consult herbicide labels for the most current and up-to-date precautions and restrictions. Copies, reproductions, or transcriptions of this document or its information must bear the statement "Produced and prepared by Purdue University or Ohio State University Extension Weed Science" unless approval is given by the author.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.